



FIFRA 24(c) Special Local Need Label (SLN)

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MINNESOTA

For Control of Diseases on Full-Season Potatoes Grown Under Irrigation

BRAVO WEATHER STIK®

EPA Reg. No. 66222-276

SLN No. MN-220002

Active Ingredient:

Chlorothalonil (tetrachloroisophalonitrile)..... 54.0%

Other Ingredients..... 46.0%

Total: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

This label is valid until December 31, 2027 or until otherwise amended, withdrawn, cancelled, or suspended

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- Read the “Limitation of Warranty and Liability” in the label booklet for BRAVO WEATHER STIK before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- The complete DIRECTIONS FOR USE can be found on the currently registered EPA Stamped Label.

Crop	Diseases (Pathogen)	Pint Product/A (lb ai/A)	Application Directions
Potato (full-season varieties grown under irrigation only)	Late Blight (<i>Phytophthora infestans</i>)	Low rate: ¾ (0.6)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).
	Early Blight (<i>Alternaria solani</i>)	-then-	
	Botrytis vine rot (<i>Botrytis cinerea</i>)	Higher rates: 1 to 1½ (0.75 to 1.125)	Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur:
	Black dot		

Crop	Diseases (Pathogen)	Pint Product/A (lb ai/A)	Application Directions
	<i>(Colletotrichum coccodes)</i>		<ul style="list-style-type: none"> • Vines close between rows • Late blight forecasting measures 18 disease severity values (DSV) • The crop reaches 300 P-days <p>Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.</p> <p>Bravo Weather Stik may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo Weather Stik label.</p>

Specific Use Directions: Do not apply more than 21 ½ pints Bravo Weather Stik (16 lb ai) per acre on full-season varieties of potato during each growing season. Do not apply within 7 days of harvest.

Bravo Weather Stik is a registered trademark of ADAMA

24(c) Registrant:
Makhteshim Agan of North America, Inc. d/b/a
ADAMA
3120 Highwoods Blvd., Suite 100
Raleigh, NC 27604

