



FIFRA 24(c) Special Local Need Label (SLN)

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MINNESOTA

For Control of Disease on Full-Season Potatoes Grown Under Irrigation

BRAVO ULTREX®

EPA Reg. No. 66222-277

SLN No. MN-220003

Active Ingredient:

Chlorothalonil (tetrachloroisophalonitrile)..... 82.5%

Other Ingredients..... 17.5%

Total: 100.0%

KEEP OUT OF REACH OF CHILDREN WARNING

This label is valid until December 31, 2027 or until otherwise amended, withdrawn, cancelled, or suspended

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- Read the “Limitation of Warranty and Liability” in the label booklet for BRAVO Ultrex before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- The complete DIRECTIONS FOR USE can be found on the currently registered EPA Stamped Label.

Crop	Diseases (Pathogens)	Lb Product/A (lb ai/A)	Application Directions
Potato (Full season varieties grown under irrigation only)	Late Blight (<i>Phytophthora infestans</i>)	Low rate: 0.7 (0.6)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).
	Early Blight (<i>Alternaria solani</i>)	-then-	
	Botrytis vine rot (<i>Botrytis cinerea</i>)	Higher rates: 0.9 to 1.36 (0.75 to 1.1)	Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur: <ul style="list-style-type: none"> • Vines close between rows • Late blight forecasting measures 18 disease severity values (DSV) • The crop reaches 300 P-days Increase water spray volume as canopy
	Black dot (<i>Colletotrichum coccodes</i>)		

Crop	Diseases (Pathogens)	Lb Product/A (lb ai/A)	Application Directions
			<p>density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.</p> <p>Bravo Ultrex may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo Ultrex label.</p>

Specific Use Restrictions: Do not apply more than 19.35 pounds of Bravo Ultrex (16 lb ai) per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment wash waters. DO NOT apply when weather conditions favor drift from treated areas. APPLY ONLY TO AREAS SPECIFIED ON LABEL.

Bravo Ultrex is a registered trademark of ADAMA

24(c) Registrant:
 Makhteshim Agan of North America, Inc. d/b/a
 ADAMA
 3120 Highwoods Blvd., Suite 100
 Raleigh, NC 27604

