



**FIFRA 24(c) Special Local Need Label (SLN)
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MINNESOTA**

S-METOLACHLOR	GROUP	15	HERBICIDE
---------------	-------	----	-----------

Dual Magnum® Herbicide

For Control of Weeds in: Crop Group 3 (Bulb Vegetables): Chive, Chinese; Chive, fresh leaves; Daylily, bulb; Garlic, bulb; Garlic, great headed, bulb; Leek; Lily, bulb; Onion, Beltsville Bunching; Onion, dry bulb; Onion, green; Onion, pearl; Onion, potato, bulb; Onion, Welsh, tops; Shallot, bulb; Shallot, fresh leaves

**EPA Reg. No. 100-816
EPA SLN No. MN-260002**

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2030

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**SYNGENTA'S SPECIAL CONDITIONS, RISKS OF USE AND DISCLAIMER FOR USE OF
DUAL MAGNUM HERBICIDE ON CROPS ON THIS 24(c) LABEL**

IMPORTANT- READ BEFORE USE

THESE CONDITIONS RISKS OF USE AND DISCLAIMER ARE REQUIRED BY SYNGENTA CROP PROTECTION LLC AND NOT SPECIFIED BY U.S. EPA OR THE OF STATE OF MINNESOTA

USE OF DUAL MAGNUM HERBICIDE (THE "PRODUCT") ON CROPS LISTED (THE "CROP") FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP. SYNGENTA RECOMMENDS THAT THE USER TEST THIS PRODUCT TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE.

Syngenta Crop Protection, LLC makes the Product available for use in the manner described in this Section 24(c) Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss.

The decision to use this Product in the manner described in this Section 24(c) Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors. Syngenta recommends that the user test this Product to determine its suitability for such intended use.

By purchasing the Product for use, or using the Product in the manner described in this Section 24(c) Labeling, you acknowledge and accept that, to the extent consistent with



applicable law:

- 1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- 2) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Section 24(c) labeling;
- 3) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- 4) Syngenta Crop Protection, LLC disclaim all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- 5) these conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Syngenta Crop Protection, LLC or their respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and main EPA-registered label must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered label.
- Dual Magnum will not control emerged weeds. For effective weed control, Dual Magnum must be applied to clean-tilled soil.
- The risk of crop injury increases when adjuvants (e.g., non-ionic surfactants, crop oils, etc.), nitrogen sources (e.g., AMS, UAN), fertilizers or other pesticides are applied with Dual Magnum.

CHIVE, Chinese, fresh leaves; CHIVE, fresh leaves; DAYLILY, bulb; GARLIC, bulb; GARLIC, great headed, bulb; LEEK; LILY, bulb; ONION, Beltsville Bunching; ONION, green; ONION, pearl; ONION, potato, bulb; ONION, Welsh, tops; SHALLOT, bulb; SHALLOT, fresh leaves

- Postemergence

Apply a broadcast application of Dual Magnum at 0.67 -1.3 pt/A (0.64-1.24 lb ai/A) postemergence starting at the two true-leaf stage of the crop. Within that rate range, use lower rates on soils relatively coarse-textured and higher rates on fine-textured soils. A band application may also be used, applying proportionally less spray mixture on the area actually treated. Dual Magnum will



not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.

Precautions: (1) In general, the risk of crop injury is greater on lighter textured soils and with higher application rates. (2) The addition of another registered herbicide as a tank mixture with Dual Magnum, will increase the risk of crop injury.

Restrictions: (1) Make only one application and apply not more than 1.3 pt/A (1.24 lb ai/A) per crop per year. Do not apply more than 1.24 lb ai/A per year of S-metolachlor-containing products to bulb vegetables. (2) Do not harvest within 21 days of the Dual Magnum application.

ONION, dry bulb

- Postemergence
-

Apply Dual Magnum at 0.67 - 1.3 pt/A (0.64-1.24 lb ai/A) broadcast starting at the two true-leaf stage of onions. Use lower rates on coarse-textured soils and higher rates on fine-textured soils. One additional application may be applied 21 or more days after the first application, if needed. A band application may also be used, applying proportionally less spray mixture on the area treated. Dual Magnum will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.

Precautions: (1) In general, the risk of crop injury is greater on lighter textured soils and with higher application rates. (2) The use of a spray adjuvant, or the addition of another registered herbicide as a tank mixture with Dual Magnum, will increase the risk of crop injury.

Restrictions: (1) Make no more than two applications of Dual Magnum per crop. (2) Apply not more than 1.3 pt/A (1.24 lb ai/A) in a single application and not more than 2.6 pt/A (2.48 lb ai/A) per year. Do not apply more than 2.48 lb ai/A per year of S-metolachlor-containing products to dry bulb onions. (3) The Dual Magnum applications cannot be less than 21 days apart. (4) Do not harvest onions within 60 days of the Dual Magnum application. (5) The harvest of green onions for and the grazing of animals on green forage or stubble are prohibited.

©2025 Syngenta

Dual Magnum® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: MN0816662AA0426