



**FIFRA 24(c) Special Local Need Label (SLN)
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MINNESOTA**

S-METOLACHLOR	GROUP	15	HERBICIDE
---------------	-------	-----------	-----------

**Dual Magnum® Herbicide
For Control of Weeds in: Crop Group 5: Broccoli; Brussels Sprouts; Cabbage;
Cauliflower; Cabbage, Chinese (Bok Choy)**

**EPA Reg. No. 100-816
EPA SLN No. MN-260004**

**This label expires and must not be distributed or used in accordance with this SLN
registration after December 31, 2030**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**SYNGENTA'S SPECIAL CONDITIONS, RISKS OF USE AND DISCLAIMER FOR USE OF
DUAL MAGNUM HERBICIDE ON CROPS ON THIS 24(c) LABEL**

IMPORTANT- READ BEFORE USE

**THESE CONDITIONS RISKS OF USE AND DISCLAIMER ARE REQUIRED BY SYNGENTA
CROP PROTECTION LLC AND NOT SPECIFIED BY U.S. EPA OR THE OF STATE OF
MINNESOTA**

**USE OF DUAL MAGNUM HERBICIDE (THE "PRODUCT") ON CROPS LISTED (THE "CROP")
FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD
REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND
UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED
BEFORE USING THE PRODUCT ON THE CROP. SYNGENTA RECOMMENDS THAT THE
USER TEST THIS PRODUCT TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE.**

**Syngenta Crop Protection, LLC makes the Product available for use in the manner
described in this Section 24(c) Labeling on the basis that, in the sole opinion of the user,
the benefits and utility derived from the use of the Product on the Crop outweigh the
potential risk of Crop injury, Crop yield reduction or Crop loss.**

**The decision to use this Product in the manner described in this Section 24(c) Labeling
must be made by each individual user on the basis of anticipated benefits versus (i) the
potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the
target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any
other relevant factors. Syngenta recommends that the user test this Product to determine
its suitability for such intended use.**

**By purchasing the Product for use, or using the Product in the manner described in this
Section 24(c) Labeling, you acknowledge and accept that, to the extent consistent with
applicable law:**



- 1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- 2) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Section 24(c) labeling;
- 3) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- 4) Syngenta Crop Protection, LLC disclaim all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- 5) these conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Syngenta Crop Protection, LLC or their respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and main EPA-registered label must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered label.
- Dual Magnum will not control emerged weeds. For effective weed control, Dual Magnum must be applied to clean-tilled soil.
- The risk of crop injury increases when adjuvants (e.g., non-ionic surfactants, crop oils, etc.), nitrogen sources (e.g., AMS, UAN), fertilizers or other pesticides are applied with Dual Magnum.

BROCCOLI, *transplanted*; **BRUSSEL SPROUTS**, *transplanted*; **CABBAGE**, *transplanted*;
CAULIFLOWER, *transplanted*; and **CABBAGE, CHINESE (BOY CHOY)**, *transplanted*
○ Preplant, prior to transplanting, or Postemergence, after transplanting

Apply Dual Magnum at a broadcast rate of 0.5 - 1.3 pt/A (0.48-1.24 lb ai/A) preplant, non-incorporated, prior to transplanting or broadcast postemergence application within 48 hours after transplanting. Use lower rates on coarse textured soils and higher rates on fine textured and muck soils. Dual Magnum will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.

Precautions: (1) In general, the risk of crop injury from the use of Dual Magnum is less with post-transplant applications than from pre-transplant applications, and the risk of crop injury is less with post-directed than from post over-the-top post-transplanting applications. (2) The application



of Dual Magnum prior to bed formation may result in crop injury due to the incorporation and/or concentration of Dual Magnum near the transplanted crop's root system. (4) Muck soils require the higher use rates; however, weed control may be reduced on muck soil. (5) The addition of another registered herbicide, especially the herbicide Goal®, will increase the risk of crop injury from postemergence applications. (6) Chinese varieties are more sensitive to injury from Dual Magnum.

Restrictions: (1) Make only one application of Dual Magnum per crop. (2) Do not apply more than 1.3 pt/A (1.24 lb ai/A) of Dual Magnum per year. Do not apply more than 1.24 lb ai/A per year of S-metolachlor-containing products to the above listed crops. (3) Do not mechanically incorporate Dual Magnum before transplanting. (4) Do not harvest these crops within 60 days of the Dual Magnum application.

©2025 Syngenta

Dual Magnum® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: MN0816711AA0426