



**FIFRA 24(c) Special Local Need Label (SLN)
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MINNESOTA**

**Reflex® Herbicide
For Control of Weeds in Pumpkin, Summer Squash, and Winter Squash**

**EPA Reg. No. 100-993
EPA SLN No. MN-260011**

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2030

**SYNGENTA'S SPECIAL CONDITIONS, RISKS OF USE AND DISCLAIMER FOR USE OF
REFLEX HERBICIDE ON CROPS ON THIS 24(c) LABEL**

IMPORTANT- READ BEFORE USE

THESE CONDITIONS RISKS OF USE AND DISCLAIMER ARE REQUIRED BY SYNGENTA CROP PROTECTION LLC AND NOT SPECIFIED BY U.S. EPA OR THE OF STATE OF MINNESOTA

USE OF REFLEX HERBICIDE (THE "PRODUCT") ON CROPS LISTED (THE "CROP") FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP. SYNGENTA RECOMMENDS THAT THE USER TEST THIS PRODUCT TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE.

Syngenta Crop Protection, LLC makes the Product available for use in the manner described in this Section 24(c) Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss.

The decision to use this Product in the manner described in this Section 24(c) Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors. Syngenta recommends that the user test this Product to determine its suitability for such intended use.

By purchasing the Product for use, or using the Product in the manner described in this Section 24(c) Labeling, you acknowledge and accept that, to the extent consistent with applicable law:

- 1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;**
- 2) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Section 24(c) labeling;**
- 3) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or**



representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;

- 4) Syngenta Crop Protection, LLC disclaim all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- 5) these conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Syngenta Crop Protection, LLC or their respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Pumpkin

Note: Pumpkin varieties may vary in their response to Reflex Herbicide. If soils are cold and wet during pumpkin germination and emergence, the use of Reflex Herbicide may delay maturity and/or reduce yields. If heavy rain occurs shortly after transplanting, Reflex Herbicide may delay maturity and or reduce yields. When using Reflex Herbicide for the first time on a particular variety, always determine crop tolerance before using. Plant only high quality seed or healthy transplants and do not plant when unfavorable growing conditions exist.

Reflex Herbicide rates below 1 pt/A (0.25 lb ai/A) are not intended to be used as a stand-alone weed control program and should be used with other herbicides and/or other methods of weed control that support weed resistance management. Refer to the Reflex Herbicide federal label for a complete list of weeds controlled.

Broadcast Application for Directed Seeded Pumpkin

Apply Reflex Herbicide at 0.5-1 pt/A (0.125-0.25 lb ai/A) as a preemergence broadcast application to direct seeded pumpkin. An overhead irrigation or rainfall event between Reflex Herbicide application and crop emergence will ensure herbicide activation and will likely reduce the potential for crop injury due to splashing.

Broadcast Application for Transplanted Pumpkin

Apply Reflex Herbicide at 0.5-1 pt/A (0.125-0.25 lb ai/A) as a pre-transplant non-incorporated preemergence broadcast application up to 7 days prior to transplanting pumpkins. Effectiveness may be reduced if untreated soil is exposed during transplanting or other cultural operations.



Avoid cultural practices that may concentrate Reflex Herbicide treated soil around the transplant root ball. An overhead irrigation or rainfall event between Reflex Herbicide application and transplanting will ensure herbicide activation and will likely reduce the potential for crop injury due to splashing.

Row Middle Application

Apply Reflex Herbicide at 0.5-1 pt/A (0.125-0.25 lb ai/A) as a row middle application after seeding but before crop emergence or prior to transplanting. For row middle application, reduce rate in proportion to area actually treated.

General Restrictions – Pumpkin

- A maximum of 1 pint of Reflex Herbicide (**or a maximum of 0.25 lb ai/A of fomesafen from any product containing fomesafen**) may be applied per acre in ALTERNATE years in all areas south of Interstate 94 in Minnesota.
- DO NOT exceed 1 pint per acre (0.25 lb ai/A) of Reflex Herbicide on pumpkin per season in all areas south of Interstate 94 in Minnesota. A maximum of 0.75 pint of Reflex Herbicide (**or a maximum of 0.1875 lb ai/A of fomesafen from any product containing fomesafen**) may be applied per acre in ALTERNATE years in all areas north of Interstate 94 and south of U.S. Highway 2, plus Betrami, Clearwater, Lake of the Woods, Kittson, Marshall, Pennington, Polk, Red Lake, and Roseau in Minnesota.
- DO NOT exceed 0.75 pint per acre (0.1875 lb ai/A) of Reflex Herbicide on pumpkin per season in all areas north of Interstate 94 and south of U.S. Highway 2, plus Betrami, Clearwater, Lake of the Woods, Kittson, Marshall, Pennington, Polk, Red Lake, and Roseau in Minnesota.
- Do not harvest pumpkin within 32 days of Reflex Herbicide application.

Summer Squash and Winter Squash

Note: Summer squash and winter squash varieties may vary in their response to Reflex Herbicide. If soils are cold and wet during germination and emergence, the use of Reflex Herbicide may delay maturity and/or reduce yields. If heavy rain occurs shortly after transplanting, Reflex Herbicide may delay maturity and or reduce yields. When using Reflex Herbicide for the first time on a particular variety, always determine crop tolerance before using. Plant only high quality seed or healthy transplants and do not plant when unfavorable growing conditions exist.

Note: Butternut squash is very sensitive to Reflex Herbicide. Do not use Reflex on butternut squash if you cannot accept some stand thinning and/or yield reduction.

Reflex Herbicide rates below 1 pt/A (0.25 lb ai/A) are not intended to be used as a stand-alone weed control program and should be used with other herbicides and/or other methods of weed control that support weed resistance management. Refer to the Reflex Herbicide federal label for a complete list of weeds controlled.

Broadcast Application for Directed Seeded Summer Squash and Winter Squash

Apply Reflex Herbicide at 0.5-1 pt/A (0.125-0.25 lb ai/A) as a preemergence broadcast application to direct seeded summer squash and winter squash. An overhead irrigation or rainfall event between Reflex Herbicide application and crop emergence will ensure herbicide activation and will likely reduce the potential for crop injury due to splashing.



Broadcast Application for Transplanted Summer Squash and Winter Squash

Apply Reflex Herbicide at 0.5-1.0 pt/A (0.125-0.25 lb ai/A) as a pre-transplant non-incorporated preemergence broadcast application up to 7 days prior to transplanting summer squash and winter squash. Effectiveness may be reduced if untreated soil is exposed during transplanting or other cultural operations. Avoid cultural practices that may concentrate Reflex Herbicide treated soil around the transplant root ball. An overhead irrigation or rainfall event between Reflex Herbicide application and transplanting will ensure herbicide activation and will likely reduce the potential for crop injury due to splashing.

Row Middle Application

Apply Reflex Herbicide at 0.5-1.0 pt/A (0.125-0.25 lb ai/A) as a row middle application after seeding but before crop emergence or prior to transplanting. For row middle application, reduce rate in proportion to area actually treated.

General Restrictions – Summer Squash and Winter Squash

- A maximum of 1 pint of Reflex Herbicide (**or a maximum of 0.25 lb ai/A of fomesafen from any product containing fomesafen**) may be applied per acre in ALTERNATE years in all areas south of Interstate 94 in Minnesota.
- DO NOT exceed 1 pint per acre (0.25 lb ai/A) of Reflex Herbicide on summer squash and winter squash per season in all areas south of Interstate 94 in Minnesota. A maximum of 0.75 pint of Reflex Herbicide (**or a maximum of 0.1875 lb ai/A of fomesafen from any product containing fomesafen**) may be applied per acre in ALTERNATE years in all areas north of Interstate 94 and south of U.S. Highway 2, plus Betrami, Clearwater, Lake of the Woods, Kittson, Marshall, Pennington, Polk, Red Lake, and Roseau in Minnesota.
- DO NOT exceed 0.75 pint per acre (0.1875 lb ai/A) of Reflex Herbicide on summer squash and winter squash per season in in all areas north of Interstate 94 and south of U.S. Highway 2, plus Betrami, Clearwater, Lake of the Woods, Kittson, Marshall, Pennington, Polk, Red Lake, and Roseau in Minnesota.
- Do not harvest summer squash or winter squash within 32 days of Reflex Herbicide application.

Rotational Crop Restrictions*

The following rotational crops may be planted after applying Reflex Herbicide at recommended rates.

Crop to be Planted	Minimum Re-cropping Interval (Months After Last Reflex Herbicide Application)
Cantaloupe	10
Cotton	0
Cucumber	10
Dry bean	0
Eggplant (transplanted only)	12
Field Corn	10
Lima Bean	4
Pea (succulent)	0
Peanuts	10
Pepper (transplanted only)	0



Crop to be Planted	Minimum Re-cropping Interval (Months After Last Reflex Herbicide Application)
Popcorn	10
Potato	0
Pumpkin	0
Rice	10
Seed corn	10
Small grains such as wheat, barley, rye	4
Snap bean	0
Soybean	0
Squash	0
Succulent soybean (edamame)	0
Succulent beans (other than snap bean and lima bean)	12
Sunflower	12
Sweet corn	10
Tomato (transplanted only)	0
Watermelon	10
All other crops not listed above	18

*These rotational crop restrictions are specific to the rates and crops specified in this label and apply only to the state of Minnesota.

Do not graze rotated small grain crops or harvest forage or straw for livestock.

©2025 Syngenta

Reflex® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: MN0993171AA1225