



# Follow the 5R's to Prevent Veterinary Drug Residue in Dairy Animals



#### **RELATIONSHIPS**

### Develop good relationships with people involved in the process

- ➤ Establish a good Veterinarian-Client-Patient Relationship (VCPR)
- ➤ Review veterinary recommendations with employees/family members who work on the farm
- ➤ Provide employees/family members with regular training on the prevention of milk residues as well as farm protocols for handling animals that have been treated



## RESPONSIBLE USE: Use and handle veterinary drugs responsibly

- Minimize use of veterinary drugs to times when they are medically necessary (less use = fewer chances for a residue)
- ➤ Store veterinary drugs for lactating and nonlactating animals separately to prevent mix-ups
- ➤ Store medicated feeds in a way that would prevent accidental use
- ➤ Properly label and store over the counter, prescription, and extra-label drugs, including information with appropriate milk and meat withdrawal times
- ➤ Develop animal treatment protocols with the help of the farm veterinarian

### **DID YOU KNOW?**

There are several ways to identify treated dairy animals such as:

- ➤ Using paint to mark an animal that has been treated where it can be easily seen by milking staff (hind leg, udder or tail)
- Using two methods to mark an animal that has been treated such as leg bands AND paint
  - Moving treated animals to a separate pen



#### **RECORDKEEPING**

#### Maintain good records to document treatments

- ➤ Use a good system to identify individual animals
- Maintain a recordkeeping system to document all treatments given
- ➤ Identify the animal before it is treated
- > Record the treatment before it is administered
- ➤ Keep treatment records for at least THREE years
- ➤ Establish a system for identifying and separating treated animals



#### **RESPECT**

#### Withdrawal times and usage limitations

- ➤ Use only veterinary drugs that are approved by the FDA for use in the species and animal class you are treating
- ➤ Use the drug only as the FDA label specifies UNLESS your veterinarian prescribes the drug for extra-label usage
- ➤ Follow withdrawal periods set by the drug manufacturers and your veterinarian (if using extra-label)



#### **REMOVE DOUBT**

- ➤ Test milk from treated, fresh and newly purchased cows for drug residues BEFORE commingling into the bulk tank
- ➤ Test bulk tank prior to milk leaving the farm, every day, every time
- ➤ Review treatment records prior to selling an animal or her milk

For additional information visit www.mda.state.mn.us/residue-prevention