

**For distribution and use in Michigan, Minnesota (for sugar beet acreage planted only for Minn-Dak Farmers Cooperative or Southern Minnesota Beet Sugar Cooperative in the following counties: Big Stone, Brown, Chippewa, Clay, Cottonwood, Douglas, Grant, Kandiyohi, Lac Qui Parle, Lyon, McLeod, Meeker, Nicollet, Otter Tail, Pope, Redwood, Renville, Sibley, Stearns, Stevens, Swift, Traverse, Watonwan, Wilkin, and Yellow Medicine), and North Dakota (for sugar beet acreage planted only for Minn-Dak Farmers Cooperative in Cass and Richland counties) under the Emergency Exemption Use Directions pursuant to Section 18 of FIFRA, as amended.**

# Provysol™

**Mefentrifluconazole Group 3 Fungicide**

## Fungicide

For control of Cercospora leaf spot (*Cercospora beticola*) in Sugar Beet

This Emergency Exemption is effective from May 31, 2019 until September 25, 2019

EPA Est. No. 51036-GA-001

EPA File Symbol Nos. 19-MI-02  
19-MN-01  
19-ND-01

**Active Ingredient\*:**

mefentrifluconazole: 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazole-1-yl)propan-2-ol. . . . . 34.93%

**Other Ingredients:** . . . . . 65.07%

**Total:** . . . . . 100.00%

\* Provysol™ fungicide contains 3.34 lbs mefentrifluconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty** on individual containers.

**In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).**

**Unused product must be returned to distributor or BASF Corporation after September 25, 2019.**

81133872 NVA 2019-05-548-0183

**Net Contents: 2 x 2.5 gallons**

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709

Product of Japan.  
Formulated in the US with domestic and imported materials.



For distribution and use in Michigan, Minnesota (for sugar beet acreage planted only for Minn-Dak Farmers Cooperative or Southern Minnesota Beet Sugar Cooperative in the following counties: Big Stone, Brown, Chippewa, Clay, Cottonwood, Douglas, Grant, Kandiyohi, Lac Qui Parle, Lyon, McLeod, Meeker, Nicollet, Otter Tail, Pope, Redwood, Renville, Sibley, Stearns, Stevens, Swift, Traverse, Watonwan, Wilkin, and Yellow Medicine), and North Dakota (for sugar beet acreage planted only for Minn-Dak Farmers Cooperative in Cass and Richland counties) under the Emergency Exemption Use Directions pursuant to Section 18 of FIFRA, as amended.

Mefentrifluconazole	Group	<b>3</b>	Fungicide
---------------------	-------	----------	-----------

# Provysol™

## Fungicide

For control of *Cercospora* leaf spot (*Cercospora beticola*) in Sugar Beet

This Emergency Exemption is effective from May 31, 2019 until September 25, 2019

EPA Est. No. 51036-GA-001

EPA File Symbol Nos. 19-MI-02  
19-MN-01  
19-ND-01

**Active Ingredient\*:**

mefentrifluconazole: 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazole-1-yl)propan-2-ol	34.93%
<b>Other Ingredients:</b>	65.07%
<b>Total:</b>	100.00%

\* Provysol™ fungicide contains 3.34 lbs mefentrifluconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty.**

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Unused product must be returned to distributor or BASF Corporation after September 25, 2019.

**Net Contents: 2.5 Gallons**

Product of U.S.A.

BASF Corporation  
26 Davis Drive, Research Triangle Park, NC 27709

NVA 2019-04-548-0020



We create chemistry

## FIRST AID

<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• <b>DO NOT</b> give anything to an unconscious person.</li></ul>
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

## Precautionary Statements

### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Harmful if inhaled. Avoid breathing spray mist. Causes moderate eye irritation. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, or viton  $\geq 14$  mils)
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove clothing/PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

**DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

This product may impact surface water quality because of runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater.

### Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### Surface Water Advisory

This product is classified as having high potential for reaching aquatic sediment via runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative

buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this active ingredient or its degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding application when rainfall is forecast to occur within 48 hours.

Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

## Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the user's possession during application. Read the entire **Directions For Use** and **Conditions of Sale and Warranty** before using this product.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

**EXCEPTION:** If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves (made of any waterproof material)
- Shoes plus socks

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

### Pesticide Storage

Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed.

### Pesticide Disposal

Wastes resulting from using this product may be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

### Container Handling

**Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

### Steps to take if material is released or spilled:

- In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

---

## Product Information

---

For optimum disease control, apply **Provysol™ fungicide** in a regularly scheduled protective spray program and use in a rotation program with **non-Group 3** fungicides labeled for use in sugar beet.

### Mode of Action

Mefentrifluconazole, the active ingredient in **Provysol™ fungicide**, inhibits the demethylation step of sterol biosynthesis (DMI), which disrupts cell membrane synthesis and is classified by the Fungicide Resistance Action Committee (FRAC) as a **Group 3** fungicide.

---

## Resistance Management

---

For resistance management, **Provysol™ fungicide** contains a **Group 3** fungicide. Any fungal population may contain individuals naturally resistant to **Provysol™ fungicide** and other **Group 3** fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

### To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Provysol™ fungicide** or other **Group 3** fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treatment area for lack of biological efficacy that might indicate possible resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or **Integrated Pest Management (IPM)** recommendations for specific crops and pathogens.
- For further information or to report suspected resistance consult your local BASF representative, extension specialist, or certified crop advisor.

---

## Application Instructions

---

- Thorough and uniform coverage is required for optimum performance.
- **Provysol™ fungicide** can be applied by ground or air.
- Application equipment must be cleaned thoroughly before and after applying this product, particularly if a product with the potential to injure crops was used before application of **Provysol™ fungicide**. Flush system with clean water.
- **Provysol™ fungicide** must be applied with other **non-Group 3** fungicides.

### Ground Application

- Use a minimum of 10 gallons of water per acre to ensure uniform and thorough canopy penetration and coverage of foliage.

**Provysol™ fungicide** label rates specify the amount of product to apply uniformly over an acre of ground on a broadcast basis.

### Aerial Application

- **Minimum spray volume per acre:** 2 gallons of spray solution per acre, unless otherwise specified in the **Detailed Use Directions** section.
- **DO NOT** apply in spray solutions less than 50% water by volume. Reduced spray volumes used in aerial application may result in physical incompatibility, reduced disease control, or crop injury particularly when mixed with other products.

### Spray Drift Management

**DO NOT** apply when conditions favor drift beyond the intended (target) application area. Drift-reduction measures include:

1. **DO NOT** apply if wind speed is 15 mph or more. Use caution when applying if nontarget crops are downwind and wind is present. **DO NOT** apply if winds are gusty.
2. Apply only when the potential for drift to adjacent sensitive areas (e.g. bodies of water or nontarget crops) is minimal and when wind is blowing away from sensitive areas.
3. Use caution when conditions are favorable for drift (drought, high temperatures, low relative humidity).
4. **DO NOT** apply during a temperature inversion. Consult local weather services before applying if conditions are favorable for inversion.



## Tank Mixing Other Products and Additives

**It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.**

**Provysol™ fungicide** can be tank mixed with other fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives. Always follow the most restrictive label use directions. See the **Detailed Use Directions** section for additional crop-specific information.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or injury may result from mixing **Provysol™ fungicide** with other products.

To minimize the likelihood of injury, before using any tank mix previously listed, test the combination on a small portion of the crop to be treated to ensure a phytotoxic response will not occur as a result of application. However, environmental variability precludes direct and consistent projection of small area test results to future use.

When an adjuvant is used with this product, BASF recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Consult a BASF representative or local agricultural authorities for more information on use of additives or adjuvants with this product.

### Compatibility Test

Before mixing components, always perform a compatibility jar test.

1. Add components in the order listed in **Mixing Order** instructions.
  - **For 10 gallons per acre spray volume:** Start with 1 pint (2 cups) of water from the intended source at the source temperature.
  - **For other spray volumes:** Adjust rates accordingly.
  - **Dry product:** Add 2 teaspoons per pound of product per acre.
  - **Liquid product:** Add 1 teaspoon per pint of product per acre.
2. Always cap the jar and invert 10 cycles after component additions.
3. When the components have all been added to the jar, let the solution stand for 15 minutes.
4. **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface, fine particles that precipitate to the bottom, or thick (clabbered) texture. **DO NOT** use any spray solution that could clog spray nozzles.

## Mixing Order

Make sure each component is thoroughly mixed and suspended before adding tank mix partners. Except when mixing products in PVA bags, maintain constant agitation during mixing and application.

1. **Water** - Fill a thoroughly clean sprayer tank 3/4 full of clean water and begin agitation.
2. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
3. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates including **Provysol™ fungicide**, or suspo-emulsions)
  - **Containers 5 gallons or less: Shake well before adding to the tank.**
  - **Containers more than 5 gallons: Recirculate before adding to the tank.**
  - Consult a BASF representative for additional information regarding agitation and recirculation.
5. **Water-soluble products**
6. **Emulsifiable concentrates** (such as oil concentrates when applicable)
7. **Water-soluble additives** [such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when applicable]
8. **Remaining quantity of water**

---

## Crop Rotation Restrictions

---

Root and tuber vegetables (except sugar beet) must not be planted within 12 months after the last application.

Sugar beet and all other crops may be planted immediately after the last application.

## Detailed Sugar Beet Use Directions

<b>Foliar</b>	<ul style="list-style-type: none"> <li>• Apply before onset of disease and continue on 14-day intervals.</li> <li>• <b>DO NOT</b> make more than two (2) sequential applications of <b>Provysol™ fungicide</b> before alternating to a labeled <b>non-Group 3</b> fungicide.</li> <li>• For Cercospora leaf spot control, <b>Provysol™ fungicide</b> must be applied with other <b>non-Group 3</b> fungicides.</li> </ul>
---------------	---

Disease Controlled		Rate per Acre (fl ozs product)		PHI; Livestock Grazing and/or Feeding Interval (All Commodities) (days)
		per application	maximum per year	
<b>Foliar</b>	Cercospora leaf spot <i>Cercospora beticola</i>	5	10	21
<ul style="list-style-type: none"> <li>• <b>Spray Interval</b> - Apply at 14-day intervals.</li> </ul>				

### Use Restrictions

- **DO NOT** apply more than 5 fl ozs (0.13 lb mefentrifluconazole) per acre per application.
- **DO NOT** make more than 2 applications.
- **DO NOT** apply more than 10 fl ozs (0.26 lb mefentrifluconazole) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.26 lb ai/acre/year of mefentrifluconazole-containing products.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Provysol is a trademark of BASF.*

© 2019 BASF Corporation  
All rights reserved.

007969-00XXX.20190422c.**NVA 2019-04-548-0019**

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



We create chemistry